



MICHAEL J. BAKER, D.P.M.

JASON D. GRAY, D.P.M.

GREGORY W. BOAKE, D.P.M

JESSICA R TAULMAN, D.P.M

BFS BUSINESS OFFICE

P O BOX 330.
Fortville, IN 46040-0330
Tel 317.863.2556
Fax 317.203.0420

COMMUNITY FOOT & ANKLE CENTER

1221 Medical Arts Blvd.
Anderson, IN 46011
Tel 765.641.0001
Fax 765.641.0003

EAST FOOT & ANKLE CENTER

161B Washington Point Dr.
Indianapolis, IN 46229
Tel 317.898.6624
Fax 317.898.6636

FOOT & ANKLE AT WESTVIEW HOSPITAL

3520 Guion Rd., Ste 102
Indianapolis, IN 46222
Tel 317.920.3240
Fax 317.920.3243

MARION FOOT CENTER

330 N. Wabash Ave, Ste 460A
Marion, IN 46952
Tel 765.664.1413
Fax 765.965.6530

BAKER FOOT SOLUTIONS SATILLITE FOOT CLINICS

BROWNSBURG

Tel 317.920.3240
Fax 317.920.3243

GEIST FAMILY PRACTICE

Tel 317.898.6624
Fax 317.898.6636

NEW CASTLE

Tel 765.664.1413
Fax 765.965.6530


SPEEDWAY

Tel 317.920.3240
Fax 317.920.3243

Ganglion Cyst

A ganglion cyst is a mass of tissue that is filled with a jelly-like fluid. The word “ganglion” means “knot” and is used to describe the knot-like mass or lump of cells that forms below the surface of the skin.

Ganglion cysts are among the most common benign soft-tissue masses. Although they most often occur on the wrist, they also frequently develop on the foot—usually on the top, but elsewhere as well. Ganglion cysts vary in size, may get smaller and larger over time and may even disappear, only to possibly return later.



What Causes a Ganglion Cyst?

Ganglion cysts are usually caused by some type of injury. For example, a ganglion cyst might develop after something drops on the foot, if the foot was twisted while walking, or after too much stress was placed on a joint or tendon. For some patients, the trauma that triggers this condition is something they do not recall. Usually the cyst appears soon after the trauma occurs.

TREATMENT

Monitoring, but no treatment. If the cyst causes no pain and does not interfere with walking, the surgeon may decide it is best to carefully watch the cyst over a period of time.

Shoe modifications. Wearing shoes that do not rub the cyst or cause irritation may be advised. In addition, placing a pad inside the shoe may help reduce pressure against the cyst.

Aspiration and injection. This technique involves draining the fluid and then injecting a steroid medication into the mass. More than one session may be needed. Although this approach is successful in some cases, in many others the cyst returns.

Surgery. When other treatment options fail or are not appropriate, the cyst may need to be surgically removed. While the recurrence rate associated with surgery is much lower than that experienced with aspiration and injection therapy, there are nevertheless cases in which the ganglion cyst returns.

